



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.


About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

L69

3 2044 106 357 924

FJ
V44



SMITHSONIAN INSTITUTION.
UNITED STATES NATIONAL MUSEUM.

SCIENTIFIC RESULTS OF EXPLORATIONS BY THE U. S. FISH
COMMISSION STEAMER ALBATROSS.

XIV.—PLANTS COLLECTED IN 1889 AT SOCORRO
AND CLARION ISLANDS, PACIFIC OCEAN.

BY

DR. GEORGE VASEY AND J. N. ROSE,

Botanist and Assistant Botanist, Department of Agriculture.

From the Proceedings of the United States National Museum, Vol. XIII, pages 145-149.

[No. 801.]

WASHINGTON:
GOVERNMENT PRINTING OFFICE.

1890.

Post, July 1912

27024

**SCIENTIFIC RESULTS OF EXPLORATIONS BY THE U. S. FISH COM-
MISSION STEAMER ALBATROSS.**

[Published by permission of Hon. Marshall McDonald, Commissioner of Fisheries.]

**No. XVI.—PLANTS COLLECTED IN 1889 AT SOCORRO AND CLARION
ISLANDS, PACIFIC OCEAN.**

BY

**DR. GEO. VASEY AND J. N. ROSE,
Botanist and Assistant Botanist, Department of Agriculture.**

Among the various islands visited by the *Albatross* in her voyage along our Pacific coast in the year 1889 were Socorro and Clarion. A small and fragmentary collection was made from each of these islands. It is to be regretted that an arrangement had not been made for a large and careful collection, not only from these but from many other interesting points visited by this ship. We are indebted to Mr. Charles H. Townsend, the ornithologist of the expedition, for this collection, which he made in connection with his other work.

Socorro Island is the largest of the Revillagigedo group, and is about 260 miles south of Cape St. Lucas, Lower California, in latitude $18^{\circ} 43' 14''$ and longitude $110^{\circ} 54' 13''$. It is about 24 miles long by 9 miles broad. It contains a small mountain, which rises to the height of 2,000 feet. Clarion or Cloud lies to the south and west of Socorro, in latitude 18° and longitude $114^{\circ} 50'$.

The flora of these islands is doubtless tropical and similar to that of Mexico. The total number of species found on the two islands was twenty-six; eighteen are from Socorro and twelve from Clarion Island, four of which they have in common. One from Socorro has hitherto been found only in Southern Lower California. Three new species and variety are now described, one from Clarion, the others from Socorro. There are three species of Sapindaceæ from these islands, three Leguminosæ from Clarion Island, three Compositæ and two Gramineæ from Socorro, and two Euphorbiaceæ, one from each island.

These plants were collected in March, 1889.

PLANTS FROM CLARION ISLAND.

PORTULACACEÆ.

Portulaca pilosa L.

STERCULIACEÆ.

Waltheria Americana L.

ZYGOPHYLLACEÆ.

Tribulus cistoides L.

SAPINDACEÆ.

Dodonæa viscosa L.*Sapindus*, sp.

LEGUMINOSÆ.

Erythrina, sp.

A single leaf of some Leguminous plant, similar to *E. crista-galli*, but the leaflet is obovate and obtuse.

Phaseolus, sp.

The whole plant cinereous, pubescent, leaflet, oval, entire; peduncle, 6 inches long; flower, purple. It resembles *P. atropurpureus* Moc. in some respects, but the leaves are dissimilar. It seems nearest No. 183 of Parry and Palmer (from Central Mexico, 1878), having similar leaves, petiole, and flowers, but the calyx lobes are more slender.

Sophora tomentosa L.

RUBIACEÆ.

Spermacoce, sp.

A single specimen obtained.

CONVOLVULACEÆ.

Ipomœa, sp.

Glabrous throughout: leaves three lobed, cordate at base; lobes obtuse to acute or even acuminate: peduncles long, five to six flowered: calyx with long acuminate lobes: corolla large (2 to 3 inches long), purple. The material of this species is poor and can not be properly identified. It does not match anything in the National Herbarium. The calyx and corolla is very similar to *I. insularis*, of Sandwich Islands, but the leaves are quite different.

LABIATEÆ.

Teucrium Townsendii sp. nov.

Low and diffuse herb almost glabrous: lower leaves (1 inch long) oblong with cuneate base, coarsely dentate: floral leaves (5 to 9 lines long) crowded, ovate, obtuse, entire: flowers solitary in the axils of

the leaves on pedicels 2 to 3 lines long: calyx 2 to 3 lines long, deeply five-lobed into ovate lanceolate divisions: corolla, lilac, 10 lines long, longer than the leaves, the lower lobes 5 lines broadly oval: seeds scarcely roughened, glabrous.

This species is very distinct from any North American representative of this genus. It resembles *T. laciniata* in its general habit, but its leaves are very different, all its floral leaves being entire and the lower ones merely dentate; the seed is also similar but smaller. *T. glandulosum* of Cedros Island has the floral leaves entire, but it is a taller form with a loose inflorescence, long pedicels and different seeds. We take pleasure in dedicating this species to the collector, Mr. Charles H. Townsend.

EUPHORBIACEÆ.

Euphorbia, sp.

PLANTS FROM SOCORRO ISLAND.

PORFULACACEÆ.

Portulaca pilosa L.

Only a single fruiting specimen.

STERCULIACEÆ.

Waltheria Americana L.

The inflorescence is unusually open and somewhat cymose.

ZYGOPHYLLACEÆ.

Tribulus cistoides L.

A very common plant on this island.

SAPINDACEÆ.

Cardiospermum Palmeri sp. nov..

Climbing over bushes; stems canescent-tomentose: leaves tomentose on both sides, biennial on petioles, half inch long; leaflets from 1 to 1½ inches long, sessile or on petiolule 3 to 4 lines long, ovate or oblong, coarsely dentate: peduncles 3 inches long: flowers rather large: the two outer sepals a half line long, broadly ovate, a little hirsute; the two inner, 2 lines long, oblong, glabrous; the two hypogynous glands short and rounded: petals white: capsule membranous, 12 to 18 lines in diameter, minutely hirsute to almost glabrous. C. H. Townsend, Socorro Island, March, 1889; Dr. Edward Palmer, La Paz, Lower California (No. 68), January 20, 1890.

Mr. Townsend got only a few capsules and a single leaf. Dr. Palmer's collection contains both flowers and fruit, and enables us to characterize this as a good species; this specimen has somewhat narrower

leaflets and more glabrous capsules, but in other respects is like the Socorro Island plant. Dr. Palmer says it is very rare in Lower California.

Two species were collected near Dr. Palmer's station by Xantus in 1859 and 1860. Neither of these was Dr. Gray able to definitely determine on account of poor material. One he refers to "*C. molle*, H. B. K. !," of which but a single specimen was collected. This we have not seen; it may possibly be the same species, but ours is certainly not *C. molle*. The other species of Xantus was referred to as "*Cardiospermum*, sp. nov." Mr. Watson thinks it may be *C. tortuosus*, although Dr. Gray was doubtful. (See Watson, Bibliographical Index, p. 79.) Bentham's plant was collected by Mr. Hind in 1837, from Magdalena Island, and Mr. Brandegee has collected it from the original station this past season. He did not find it at all on the main-land. Our plant is clearly not this species. *C. Halicacabum*, the only other species from the peninsula, is almost glabrous, with smaller flowers. Four or five other species occur in Mexico, but differ in hypogynous glands, foliage, etc.

The species to which it comes nearest is *C. Loxense*, H. B. K., from Peru, South America, but this has leaves with petiole $1\frac{1}{2}$ to 2 inches long, the outer sepal half the length of the inner, and with the capsule "pubescenti-hirtelles," etc.

Dr. Gray referred here, with some uncertainty, a plant from Wilkes's expedition, with perfectly glabrous capsules. Our plant resembles it, but has broader leaflets, longer peduncles, etc.

Dodonæa viscosa L.

RUBIACEÆ.

Spermacoce (Boneria), sp.

Species resembling *S. podocephala* Gray.

COMPOSITÆ.

Erigeron, sp.

Viguiera deltoidea Gray, var. *Townsendii*, var. nov.

Leaves entire and opposite: thinner and with less scabrosity than the type: rays five to eight.

Perityle Socorroensis Rose. Bot. Gaz., xv, 118. The type.

SOLANACEÆ.

Physalis glabra Benth. (?)

This may be a new species, but the material is insufficient and for the present it is referred to the above species. It is quite glabrous throughout: the largest leaves $1\frac{1}{2}$ inches long, cordate at base, obtuse at apex: corolla small (2 lines broad), with a large purple eye, anther bluish green.

ACANTHACEÆ.

Elytraria tridentata Vahl.

VERBENACEÆ.

Lantana involucrata L.

ARISTOLOCHIACEÆ.

Aristolochia brevipes Benth.

Probably this species, but material very scanty.

LORANTHACEÆ.

Phoradendron rubrum Griseb.

A single specimen sent without flower or fruit; the identification is uncertain; the leaves correspond very well with specimen in the National Herbarium.

EUPHORBIACEÆ.

Euphorbia, sp.

CYPERACEÆ.

Fimbristylis, sp.

GRAMINEÆ.

Cenchrus myosuroides H. B. K.*Heteropogon contortus* R. & S.

FILICES.

Cheilanthes Wrightii Hook.

“Probably this species, but material too poor for exact determination.” H. Seaton.

MARCH 5, 1889.

